

**703**

Li X<sup>1</sup>, Di B<sup>1</sup>, Zhao E<sup>1</sup>, Xu G<sup>1</sup>

1. Senggong Hosipital in Heilongjiang Province

## **THE THERAPEUTIC EFFECTS OF SOLIFENACIN SUCCINATE ON CYSTOSPASM IN PATIENTS WITH CONTINENT BLADDER STOMA**

### Hypothesis / aims of study

To investigate the therapeutic effects of solifenacin succinate on cystospasm in patients with continent bladder stoma.

### Study design, materials and methods

Twenty patients with cystospasm after establishment of continent bladder stoma received solifenacin for 4 weeks. The symptoms of cystospasm were evaluated and compared and the adverse reactions were recorded before and after treatment.

### Results

Solifenacin therapy significantly improved the pain or discomforts in patients with cystospasm; only 1 patient experienced dry mouth and related symptoms.

### Interpretation of results

Solifenacin succinate is safe and effective in treatment of cystospasm in patients with continent bladder stoma.

### Concluding message

cystospasm; solifenacin succinate; continent bladder stoma

### Disclosures

**Funding:** No **Clinical Trial:** Yes **Public Registry:** No **RCT:** Yes **Subjects:** HUMAN **Ethics not Req'd:** It is investigor driven study and there is not treatment before for these kinds of patients **Helsinki:** Yes **Informed Consent:** Yes