

470

Wei W¹, Yang J¹

1. Dalian Friendship Hospital

SOLIFENACIN COMBINED WITH BIOFEEDBACK IN THE TREATMENT OF FEMALE OVERACTIVE BLADDER

Hypothesis / aims of study

To evaluate the effect solifenacin combined with biofeedback in the treatment of female Overactive Bladder (OAB)

Study design, materials and methods

23 cases of female OAB were treated with solifenacin(28 days) combined with biofeedback(20 days). The effects were evaluated after 3 months and followed up for 3 months.

Results

The total efficiency rate was 87%the cure rate, the alleviation rate, and the nullity rate was 56.5%(13 cases), 30.4%(7 cases), 13%(3 cases)respectively. 19 cases were followed up for 6 months and no one relapsed.

Interpretation of results

solifenacin combined with biofeedback could be an effective therapeutic method for patients with OAB.

Concluding message

overactive bladder ; biofeedback ; solifenacin

References

1. Abrams P,Cardozo L,Fall M,et al.The standardization of terminology of lower urinary tract function,report from the standardization subcommittee of the international continence societyj.Neurourol Urodyn,2002.21(2):167.
2. Claudio T,Fernanda C,Fernanda Aw,et al Overactive bladder:Prevalence and implications in Brazil European Urology,2006,49:1087-1092.
3. Reese PR,Pleil AM,Okano GJ,et al.Multinational study of reliability and validity of the King'S Health Questionnaire in patients with overactive bladder.Qual Life Res,2003,12:427-442.

Disclosures

Funding: No **Clinical Trial:** Yes **Public Registry:** No **RCT:** Yes **Subjects:** HUMAN **Ethics not Req'd:** It is investigator driven study and there is not treatment before for these kinds of patients **Helsinki:** Yes **Informed Consent:** Yes