

MEDIUM-TERM FUNCTIONAL RESULTS OF LAPAROSCOPIC SACROCOLPOPEXY. A RETROSPECTIVE, SINGLE-CENTER STUDY ABOUT 81 CASES.

Hypothesis / aims of study

Laparoscopic sacrocolpopexy (LSC) offers the possibility of global repair of pelvic organ prolapse (POP) with low morbidity and a relatively easy approach to the retrocervical space when compared with laparotomy [1, 2]. The available data on the functional results of LSC are not sufficient at the present time [3].

The aim of the study was to analyze the medium-term functional results of surgical repair of POP by LSC.

Study design, materials and methods

81 patients operated on between October 2003 and December 2008 were included in the study. Intra- and postoperative data were collected. A self questionnaire of symptoms and quality of life was sent to all patients. Responses from 62 patients were collected and analyzed.

Results

The mean age was 54 years (range 36 – 72 years), and the mean follow-up time was 18 months (median 12.5). Of the patients who responded, 62% were menopausal, 3% had a past history of a total hysterectomy and 8% had recurrent prolapse. All the patients suffered from a POP stage \geq II, uni- or multi-compartmental. In addition, 15% suffered from stress urinary incontinence (SUI), and 40% had a positive ALPP with reduced prolapsus on urodynamic study (UDS). The meshes used (71% of cases with multifilament polyester) were fixed by nonabsorbable stitches in 69% of the cases or by Tackers in 31% of the cases. Fixation on the levator ani muscle was carried out in 80% of the cases. Subtotal hysterectomy was done in 6% of the cases, and treatment of SUI by a suburethral sling was carried out in 23% of the cases. The mean time of the intervention was 184 minutes. In spite of one case of laparoconversion due to hypercapnia, no severe intraoperative complications were found. The average hospital stay was 3.5 days. The postoperative global satisfaction was 93.4%. The rate of de novo constipation was 27.8%, and the rate of de novo SUI was 11.4% (of which 1.6% were reoperated). One patient suffered from de novo dyspareunia (1.6%). No cases of vaginal erosion were found. The rate of anatomical success was 96.3%. In fact, we regret that 3 cases of anterior compartment recurrence occurred (at 1, 8 and 12 months), requiring a second intervention via the vagina (2) or by laparoscopy (1). 94% of patients stated that they were completely cured, and 91% said that they would recommend this intervention to a friend.

Interpretation of results

According to this study, the medium-term anatomical and functional results were excellent. Severe constipation was a minor, but frequent, postoperative complication. LSC seems to give both less postoperative pain and less dyspareunia than transvaginal POP repair. Studies comparing laparoscopic and transvaginal methods are needed to confirm this proposal.

Concluding message

LSC offers the possibility of global repair of POP with low morbidity and a high satisfaction rate. The technique requires perfect knowledge of pelvic anatomy and experienced surgeons.

References

1. The current status of laparoscopic sacrocolpopexy: a review. Ganatra AM, Rozet F, Sanchez-Salas R, Barret E, Galiano M, Cathelineau X, Vallancien G. Eur Urol. 2009 May;55(5):1089-103. Epub 2009 Feb 4.
2. Twenty years of laparoscopic sacrocolpopexy: where are we now? Gabriel B, Nassif J, Barata S, Wattiez A. Int Urogynecol J. 2011 Sep;22(9):1165-9. Epub 2011 Mar 18.
3. Surgical management of pelvic organ prolapse in women: the updated summary version Cochrane review. Maher CM, Feiner B, Baessler K, Glazener CM. Int Urogynecol J. 2011 Nov;22(11):1445-57. Epub 2011 Sep 17.

Disclosures

Funding: NO SOURCE OF FUNDING **Clinical Trial:** Yes **Public Registry:** No **RCT:** No **Subjects:** HUMAN **Ethics not Req'd:** RETROSPECTIVE STUDY. PATIENTS CONSENT OBTAINED **Helsinki:** Yes **Informed Consent:** Yes